

PRODUCTION OF POLYMER-PRECURSOR CARBOXYLIC ACIDS FROM MUNICIPAL SOLID WASTE: FERMENTATION, CHAIN ELONGATION, AND DOWNSTREAM PROCESSING

SAFOORA MIRMOHAMADSADEGHI^{1*} AND KEIKHOSRO KARIMI²

¹Department of Biotechnology, Faculty of Biological Science and Technology, University of Isfahan, Isfahan 81746-73441, Iran


²Department of Chemical Engineering, Vrije Universiteit Brussel, 1050 Brussels, Belgium

*Corresponding author: s.mirmohamadsadeghi@bio.ui.ac.ir

HIGHLIGHTS

- MSW can be converted into polymer-precursor carboxylic acids.
- Butyrate, caproate, lactate, and succinate pathways are briefly reviewed.
- Lactate and butyrate appear to be more economically promising than succinate.
- Key challenges include contaminants, process integration, and separation.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article History:

Received: 5 June 2026

Revised: 24 June 2026

Accepted: 25 June 2026

Published: 15 July 2026

Keywords:

Municipal solid waste, chain elongation, carboxylate platform, butyric acid, caproic acid, lactic acid, succinic acid, biopolymer precursors, waste valorisation

ABSTRACT

Municipal solid waste (MSW) represents an underutilised feedstock for the production of polymer-precursor carboxylic acids via the carboxylate platform and chain elongation. This preliminary review examines the production of four key building blocks from the organic fraction of MSW: Butyric acid (C4, precursor to polyhydroxybutyrate), caproic acid (C6, precursor to polyhydroxycaproate), lactic acid (C3, precursor to polylactic acid), and succinic acid (C4, precursor to polybutylene succinate). For each chemical, fermentation strategies, key microorganisms, and separation technologies are reviewed. Lactic acid achieves the highest yields and titres but suffers from optical purity issues from mixed MSW

cultures. Butyric acid offers the best overall balance of yield, productivity, and separation feasibility. Caproic acid requires chain elongation with external electron donors, resulting in lower yields and titres, and limiting its economic viability. Succinic acid is the least mature, with low yields and high separation costs. Key knowledge gaps include contaminant impacts on chain-elongating consortia, process integration at pilot scale, and low-cost separation technologies compatible with MSW solids.

© UMT Press

Introduction

The accumulation of Municipal Solid Waste (MSW) presents one of the most pressing environmental challenges of the 21st century, with global annual generation exceeding 2 billion tonnes, of which less than 20% is effectively recycled or valorised (Kaza *et al.*, 2018). Simultaneously, the petrochemical industry's reliance on fossil resources for polymer production has driven interest in bio-based alternatives (Zhu *et al.*, 2016). This review bridges these two issues by examining the production of polymer precursor carboxylic acids from the Organic Fraction of

MSW (OFMSW) via the carboxylate platform and chain elongation.

The most important chemicals in this context are short-chain volatile fatty acids (VFAs; C2–C5) and medium-chain carboxylates (MCCs; C6–C8), some of which act as direct monomers, while others serve as carbon sources for the biosynthesis of bioplastics and other bio-based polymers (Angenent *et al.*, 2016). Table 1 summarises the key chemicals, their properties, and polymer applications.

Table 1: Key polymer precursor carboxylic acids from MSW fermentation

Chemical	Formula	Chain Length	Polymer Application	Polymer Precursor Role	References
Butyric acid	C ₄ H ₈ O ₂	C4	Polyhydroxybutyrate (PHB)	Carbon source for PHB production	Vlaeminck <i>et al.</i> (2022)
Caproic acid	C ₆ H ₁₂ O ₂	C6	Polyhydroxyalkanoates (PHAs)	Carbon source for the production of 3HHx-containing PHAs	Hahn <i>et al.</i> (2024)
Lactic acid	C ₃ H ₆ O ₃	C3	Polylactic acid (PLA)	Direct monomer for PLA	Swetha <i>et al.</i> (2023)
Succinic acid	C ₄ H ₆ O ₄	C4	Polybutylene succinate (PBS)	Direct monomer for PBS	Rajendran and Han (2023)

While lactic and succinic acids are primarily produced via classical fermentation pathways (Beauprez *et al.*, 2010), caproic acid and, under certain conditions, butyric acid, can also be generated through chain elongation, a reverse β -oxidation process in which shorter-chain carboxylates (e.g., acetate) are elongated using electron donors such as ethanol or lactate (Angenent *et al.*, 2016). MSW provides an abundant, low-cost feedstock for these processes, but its heterogeneity, contaminants, and variable

composition pose substantial challenges (Campuzano and González-Martínez, 2016). Figure 1 presents a conceptual overview of the major biological conversion routes leading to polymer precursor production from MSW. Accordingly, this review examines the production of each key polymer precursor from MSW, covering fermentation strategies, key microorganisms, and separation technologies, and concludes with a comparative analysis of technical readiness and economic viability




Figure 1: Overall process routes for production of polymer precursor carboxylic acids from MSW

Butyric Acid (C4) Production from MSW

Butyric acid is a four-carbon saturated fatty acid that can serve as a carbon substrate for microbial polyhydroxybutyrate (PHB) biosynthesis, while PHB itself is a

biodegradable thermoplastic composed of 3-hydroxybutyrate monomer units and exhibits properties similar to polypropylene (Vlaeminck *et al.*, 2022).

Fermentation

Butyric acid production from MSW typically employs anaerobic fermentation of the OFMSW (Strazzera *et al.*, 2018). Two main approaches exist: Single-stage fermentation, where OFMSW is converted to butyric acid in a single stage, and phase-separated systems in which hydrolysis and acidogenesis are operated in separate reactors to improve process control and butyric acid selectivity (Holl *et al.*, 2022).

Operational parameters strongly influence butyric acid production during the acidogenic fermentation of OFMSW. Previous studies have shown that mildly acidic conditions (approximately pH 5.5 to 6.5) favour the accumulation of butyric acid, while both mesophilic and thermophilic temperatures have been successfully applied. However, the optimal hydraulic retention time, organic loading rate, and butyric acid productivity vary considerably depending on feedstock characteristics, reactor configuration, and microbial community composition (Atasoy *et al.*, 2018; Strazzera *et al.*, 2018).

Temperature has also been shown to influence product selectivity in chain-elongating mixed cultures. Recent evidence suggests that decreasing the operating temperature may shift the fermentation profile from caproic acid toward butyric acid production by altering microbial community composition, with lower temperatures

favouring the enrichment of butyrate-producing populations such as *Clostridium sensu stricto* while reducing the abundance of caproate-producing microorganisms (Sabbe *et al.*, 2025). A major challenge remains the concurrent production of competing metabolites, particularly acetate and propionate, which may reduce butyric acid selectivity in mixed-culture fermentations (Angenent *et al.*, 2016).

Microorganisms

The microbial communities responsible for butyric acid production during OFMSW fermentation are dominated by members of the phylum *Firmicutes*, particularly the class *Clostridia* (Vital *et al.*, 2014). Among them, species belonging to the genera *Clostridium* and *Megasphaera* have frequently been associated with butyric acid formation in mixed-culture anaerobic systems (Vital *et al.*, (2014); Atasoy *et al.*, (2018). Environmental factors, especially pH, play a critical role in shaping microbial community composition and metabolic pathways. Consequently, understanding the ecological interactions within mixed microbial consortia is essential for developing strategies to enhance butyric acid selectivity during OFMSW fermentation (Atasoy *et al.*, 2018; Strazzera *et al.*, 2018). The most important butyrate-producing microorganisms, substrate preferences, and representative yields are summarised in Table 2.

Table 2: Representative butyrate-producing microorganisms, substrate preferences, and reported yields

Species	Substrate Preference	Notes	References
<i>Clostridium butyricum</i>	Glucose, starch	Most studied, robust	Fu <i>et al.</i> (2022)
<i>Clostridium acetobutylicum</i>	Multiple sugars	Also produces solvents	Jang <i>et al.</i> (2014)
<i>Megasphaera elsdenii</i>	Lactate, glucose	Lactate-utilising	Cabral and Weimer (2024)
<i>Butyrivibrio fibrisolvens</i>	Cellulose, hemicellulose	Degrades fibres	Asanuma <i>et al.</i> (2005)

Caproic Acid (C6) Production from MSW

Caproic acid (hexanoic acid) is a six-carbon medium-chain carboxylate that has attracted increasing interest as a platform chemical because of its potential use as a carbon source for the biosynthesis of medium-chain-length polyhydroxyalkanoates (mcl-PHAs) in addition to its applications in lubricant formulations, antimicrobial agents, and flavour compounds (Hahn *et al.*, 2024, Wu *et al.*, 2024). Production of caproic acid from MSW typically relies on chain elongation, in which shorter-chain carboxylates such as acetate and butyrate are elongated using electron donors including ethanol and lactate (Wang *et al.*, 2023).

Chain elongation is a microbial

process that transforms short-chain carboxylates into more valuable medium-chain carboxylates through the reverse β -oxidation pathway. During this process, electron donors provide reducing equivalents that drive the sequential elongation of carboxylate chains by two carbon atoms per cycle (Wang *et al.*, 2023). Among the products generated via reverse β -oxidation, caproic acid has received considerable attention because of its potential applications as a platform chemical and precursor for the synthesis of bio-based materials (Wu *et al.*, 2024). A simplified overview of the reverse β -oxidation pathway involved in caproate production is presented in Figure 2.




Figure 2: Reverse β-oxidation pathway for caproic acid production from short-chain carboxylates via chain elongation

Fermentation

Caproate production from MSW primarily occurs through the reverse β-oxidation pathway, which constitutes the metabolic basis of microbial chain elongation. As illustrated in Figure 2, the reverse β-oxidation cycle involves the condensation of acetyl-CoA with an acyl-CoA intermediate, followed by reduction, dehydration, and a

second reduction step. Through repeated cycles, short-chain carboxylates can be elongated by two carbon atoms per cycle, enabling the conversion of acetate to butyrate and subsequently to caproate. A simplified stoichiometric representation of caproate formation from acetate and ethanol is shown below:



In contrast to butyrate production, caproate synthesis through chain elongation is strongly influenced by electron donor availability and microbial competition. Reported caproate titres (5.4 to 14.6 g/L) and productivities from OFMSW-derived substrates vary substantially depending on feedstock composition, reactor

configuration, and operational conditions (Pu *et al.*, 2026).

The large-scale implementation of OFMSW-based caproate production is hindered by several interconnected challenges. The heterogeneous nature of food waste (FW) leads to unstable fermentation performance and frequent imbalances in pH and electron donor availability. In mixed

microbial communities, competition from lactate producers, methanogens, and other fermentative microorganisms diverts carbon and reduces power away from chain elongation, thereby reducing caproate selectivity. Moreover, process efficiency is often constrained by product inhibition, difficulties in maintaining optimal operating conditions, and the presence of inhibitory compounds such as ammonium and salts. Addressing these limitations through improved process control, microbial management, and integrated product recovery strategies will be crucial for the development of economically viable caproate biorefineries based on food waste (Pu *et al.*, 2026).

Microorganisms

Caproate production is mediated by specialised chain-elongating bacteria (CEB), primarily from the phyla Firmicutes and Proteobacteria (Wang *et al.*, 2023). Among

them, *Clostridium kluyveri* is the best-characterised caproate producer and represents an obligate chain elongator that relies on ethanol and short-chain carboxylates rather than sugars for growth (Fernández-Blanco *et al.*, 2024). In MSW consortia, *C. kluyveri* often coexists with *C. acetobutylicum* (which produces butyrate) and *Lactobacillus spp.* (which produce lactate) (Ulčar *et al.*, 2024).

A key recent discovery is lactate-mediated chain elongation, where lactate replaces ethanol as the electron donor, reducing external chemical addition costs (Wang *et al.*, 2022). However, maintaining stable lactate-driven chain elongation in mixed cultures remains challenging because of complex microbial interactions and competing metabolic pathways (Wu *et al.*, 2023).

Caproic acid recovery is more challenging than butyric acid due to its higher hydrophobicity and lower concentration in fermentation broth (Wang *et al.*, 2023).

Table 3: Representative chain-elongating bacteria for caproate production, electron donors, and reported yields

Species	Electron Donor	C4 Intermediate	References
<i>Clostridium kluyveri</i>	Ethanol + acetate	Butyrate (trace)	Reddy <i>et al.</i> (2018)
<i>Megasphaera elsdenii</i>	Lactate + acetate	Butyrate (major)	Cabral and Weimer (2024)
<i>Eubacterium pyruvativorans</i>	Ethanol + acetate	Butyrate (minor)	Hernandez-Correa <i>et al.</i> (2023)
<i>Ruminococcaceae</i> bacterium CPB6	Ethanol + acetate	Butyrate (intermediate)	Zhu <i>et al.</i> (2017)

Other Polymer Precursors from MSW

While butyric and caproic acids are the primary chain elongation products, MSW

fermentation also produces other valuable polymer precursors, including lactic acid for

polylactic acid (PLA) production (Song *et al.*, 2024) and succinic acid for polybutylene succinate (PBS) synthesis (Ioannidou *et al.*, 2023).

Lactic Acid (C3) – Fermentation

Lactic acid is mainly produced through the fermentation of carbohydrate-rich OFMSW by lactic acid bacteria, particularly species belonging to the genera *Lactobacillus* and *Weizmannia* (Song *et al.*, 2024). Process performance is strongly influenced by operational conditions such as pH and temperature, with mildly acidic conditions and mesophilic to moderately thermophilic temperatures generally favouring lactic acid accumulation (Aulitto *et al.*, 2024, Tang *et al.*, 2024).

A major challenge for PLA production is the requirement for high optical purity (i.e., a high proportion of the L-isomer), as mixed microbial cultures may produce both L- and D-lactic acid isomers (Swetha *et al.*, 2023). In this regard, *Weizmannia coagulans* (formerly *Bacillus*

coagulans) has attracted increasing attention because of its thermophilic characteristics and ability to produce optically pure L-lactic acid under industrially relevant conditions (Aulitto *et al.*, 2024).

Lactic Acid – Microorganisms

Lactic acid production from MSW-derived substrates is commonly mediated by lactic acid bacteria, with the optical purity of the final product depending strongly on the microbial strain employed. Among the available producers, *Weizmannia coagulans* (formerly *Bacillus coagulans*) has received particular attention because of its thermophilic nature, which reduces contamination risks, and its ability to produce optically pure L-lactic acid (Aulitto *et al.*, 2024). Other microorganisms commonly used for lactic acid production include *Lacticaseibacillus casei*, *Lactobacillus delbrueckii*, and *Lactiplantibacillus plantarum*, although their stereospecificity is often strain-dependent (Table 4).

Table 4: Representative lactic acid-producing microorganisms, optical isomer specificity, and reported yields

Species	Isomer	References
<i>Lactobacillus casei</i>	L	Panesar <i>et al.</i> (2010)
<i>Lactobacillus delbrueckii</i>	D	Mukherjee <i>et al.</i> (2025)
<i>Weizmannia coagulans</i> (<i>Bacillus coagulans</i>)	L (> 99%)	Pamueangmun <i>et al.</i> (2023)
<i>Lactobacillus plantarum</i>	DL	Sudhakar and Dharani (2022)

Lactic Acid – Separation

Downstream recovery of lactic acid remains a major economic challenge because of its high solubility in aqueous fermentation broths. Conventional separation strategies

include precipitation, solvent extraction, electrodialysis, and membrane-based processes. Calcium lactate precipitation is still widely used because of its operational simplicity and technological maturity,

whereas reactive extraction has emerged as a promising alternative for improving recovery efficiency and reducing downstream processing costs (Swetha *et al.*, 2023; Song *et al.*, 2024).

Succinic Acid (C4) – Fermentation

Succinic acid can be produced from OFMSW-derived hydrolysates through anaerobic fermentation by succinate-producing microorganisms (Yi *et al.*, 2013; Mitrea *et al.*, 2024). These processes are typically operated under near-neutral pH conditions and require CO₂ supplementation to support carboxylation reactions involved in succinate biosynthesis. Reported succinic acid yields and final concentrations vary considerably depending on substrate characteristics, pretreatment methods, and microbial strains employed (Yang *et al.*,

2024). A major challenge associated with this process is the need to maintain strict anaerobic conditions together with adequate CO₂ availability, which can increase operational complexity and overall production costs.

Succinic Acid – Microorganisms

Among the microorganisms investigated for bio-based succinic acid production, *Actinobacillus succinogenes* has emerged as one of the most promising candidates because of its broad substrate utilisation spectrum, including both hexose and pentose sugars, as well as its high capacity for CO₂ fixation (Yang *et al.*, 2024). These characteristics make this organism particularly attractive for the valorisation of heterogeneous MSW-derived hydrolysates.

Table 5: Representative succinic acid-producing microorganisms, yields, and fermentation characteristics

Species	Byproducts	Notes	References
<i>Actinobacillus succinogenes</i>	Acetate, formate	Most studied, high yield	Escanciano <i>et al.</i> (2024)
<i>Mannheimia succiniciproducens</i>	Acetate	CO ₂ -tolerant	Kim <i>et al.</i> (2025)
<i>Basfia succiniciproducens</i>	Acetate, lactate	Grows on glycerol	Rochón <i>et al.</i> (2024)
<i>Saccharomyces cerevisiae</i> (engineered)	Ethanol	Less robust	Malubhoy <i>et al.</i> (2022)

Succinic Acid – Separation

Succinic acid recovery from MSW fermentation broths is highly complex due to its low baseline titres and the simultaneous presence of co-produced organic acids (e.g., acetic and formic acids) which possess overlapping chemical properties. Current separation technologies, including reactive

extraction, electrodialysis, and crystallisation, suffer from severe membrane fouling and high chemical consumption caused by suspended MSW solids and colloidal impurities. Consequently, few currently available downstream methods appear economically viable for MSW-

derived succinic acid at present titres, rendering this pathway the least technologically mature of the four examined precursors (Lithourgidis *et al.*, 2023).

Critical Research Priorities

Advancing MSW-to-polymer-precursor biorefineries requires bridging fundamental microbial ecology with applied process engineering:

1. **Contaminant Resilience:** Systematic assessment of chain-elongation consortium stability under real MSW contaminant profiles (microplastics, heavy metals, xenobiotics) to enable non-sterile operation and cost reduction.
2. **Microbiome Engineering:** Development of multi-omics-guided approaches for selective enrichment of target acidogens, including synthetic consortium design and non-sterile microbiome-management protocols.
3. **Lactate-Driven Integration:** Optimisation of lactate-mediated chain elongation directly coupled to homolactic fermentation from cellulosic MSW fractions, eliminating external electron donor costs.
4. **Fouling-Resistant Separation:** Innovation in downstream technologies (membrane systems, reactive extraction, electrochemical methods) robust to high-solids MSW broths and capable of in-situ product recovery.
5. **Pilot-Scale Demonstration:** Integrated demonstration systems

combining hydrolysis, fermentation, and separation stages to validate economic performance and environmental metrics at pre-commercial scales.

Conclusions

Municipal solid waste represents a promising and underutilised feedstock for the sustainable production of polymer precursors. The technical feasibility of these pathways varies significantly by product. Lactic acid appears to be the most technologically mature option, delivering high yields and titres from OFMSW. However, the production of racemic DL-lactate from mixed MSW cultures remains a critical barrier for PLA synthesis, necessitating advanced strain engineering or the complex separation of L- and D-isomers. Meanwhile, butyric acid offers a favourable balance of yield and productivity. The integration of fermentation, chain elongation, and efficient product recovery could substantially improve the economic viability of MSW biorefineries and contribute to the transition toward a circular bioeconomy.

Chain elongation enables caproic acid production but introduces unique operational bottlenecks. Generating caproic acid from MSW requires external electron donors such as ethanol or lactate alongside longer hydraulic retention times, which ultimately results in lower yields and titres than those observed for butyric acid. While lactate-mediated chain elongation represents a promising recent development that can substantially reduce external chemical costs, its overall process stability remains inferior to conventional ethanol-based systems.

To advance MSW-to-polymer-precursor biorefineries, future research must bridge the gap between fundamental microbial ecology and process engineering. Key priorities include assessing the resilience of chain-elongation consortia against real-world MSW contaminants and developing non-sterile microbiome management strategies—potentially guided by multi-omics tools—to selectively enrich target acidogens. Furthermore, optimising self-sustaining, lactate-driven chain elongation from the cellulosic fraction of MSW remains critical to eliminate external electron donor costs. Finally, scaling up these technologies requires significant innovations in robust, fouling-resistant downstream separation systems capable of operating directly with high-solid MSW broths.

Acknowledgements

The authors would like to express their gratitude to the anonymous reviewers for their insightful comments and constructive feedback, which greatly improved the quality of this manuscript. We also thank the editorial and administrative staff for their valuable support and assistance throughout the peer-review and publication process.

Conflict of Interest Statement

The authors declare no conflict of interest.

References

Angenent, L. T., Richter, H., Buckel, W., Spirito, C. M., Steinbusch, K. J. J., Plugge, C. M., Strik, D. P. B. T. B., Grootsholten, T. I. M., Buisman, C.

J. N., & Hamelers, H. V. M. (2016). Chain elongation with reactor microbiomes: Open-culture biotechnology to produce biochemicals. *Environmental Science & Technology*, 50(6), 2796–2810. <https://doi.org/10.1021/acs.est.5b04847>

Asanuma, N., Ishiwata, M., Yoshii, T., Kikuchi, M., Nishina, Y., & Hino, T. (2005). Characterization and transcription of the genes involved in butyrate production in *Butyrivibrio fibrisolvens* type I and II strains. *Current Microbiology*, 51(2), 91–94. <https://doi.org/10.1007/s00284-005-4477-x>

Atasoy, M., Owusu-Agyeman, I., Plaza, E., & Cetecioglu, Z. (2018). Bio-based volatile fatty acid production and recovery from waste streams: Current status and future challenges. *Bioresource Technology*, 268, 773–786. <https://doi.org/10.1016/j.biortech.2018.07.042>

Aulitto, M., Alfano, A., Maresca, E., Avolio, R., Errico, M. E., Gentile, G., Cozzolino, F., Monti, M., Pirozzi, A., Donsì, F., Cimini, D., Schiraldi, C., & Contursi, P. (2024). Thermophilic biocatalysts for one-step conversion of citrus waste into lactic acid. *Applied Microbiology and Biotechnology*, 108(1), 155. <https://doi.org/10.1007/s00253-023-12904-7>

Beauprez, J. J., De Mey, M., & Soetaert, W. K. (2010). Microbial succinic acid

- production: Natural versus metabolic engineered producers. *Process Biochemistry*, 45(7), 1103–1114. <https://doi.org/10.1016/j.procbio.2010.03.035>
- Cabral, L. D. S., & Weimer, P. J. (2024). *Megasphaera elsdenii*: Its role in ruminant nutrition and its potential industrial application for organic acid biosynthesis. *Microorganisms*, 12(1), 219. <https://doi.org/10.3390/microorganisms12010219>
- Campuzano, R., & González-Martínez, S. (2016). Characteristics of the organic fraction of municipal solid waste and methane production: A review. *Waste Management*, 54, 3–12. <https://doi.org/10.1016/j.wasman.2016.05.016>
- Escanciano, I. A., Ladero, M., Blanco, A., & Santos, V. E. (2024). Succinic acid production by *Actinobacillus succinogenes* using acid and enzymatic hydrolysates of potato and beer wastes and repeated batch operation. *Biomass and Bioenergy*, 181, 107034. <https://doi.org/10.1016/j.biombioe.2023.107034>
- Fernández-Blanco, C., Veiga, M. C., & Kennes, C. (2024). Carbon dioxide as key player in chain elongation and growth of *Clostridium kluyveri*: Insights from batch and bioreactor studies. *Bioresource Technology*, 394, 130192. <https://doi.org/10.1016/j.biortech.2023.130192>
- Fu, H., Yue, Z., Feng, J., Bao, T., Yang, S. T., Cai, Y., & Wang, J. (2022). Consolidated bioprocessing for butyric acid production from raw cassava starch by a newly isolated *Clostridium butyricum* SCUT620. *Industrial Crops and Products*, 187, 115446. <https://doi.org/10.1016/j.indcrop.2022.115446>
- Hahn, T., Alzate, M. O., Leonhardt, S., Tamang, P., & Zibek, S. (2024). Current trends in medium-chain-length polyhydroxyalkanoates: Microbial production, purification, and characterization. *Engineering in Life Sciences*, 24(6), 2300211. <https://doi.org/10.1002/elsc.202300211>
- Hernandez-Correa, E., de María Cuervo-López, F., Cervantes, F. J., & Buitron, G. (2023). Efficient conversion of wine lees into medium-chain carboxylic acids by mixed culture using an in-line extraction system. *Journal of Water Process Engineering*, 55, 104263. <https://doi.org/10.1016/j.jwpe.2023.104263>
- Holl, E., Steinbrenner, J., Merkle, W., Krümpel, J., Lansing, S., Baier, U., Oechsner, H., & Lemmer, A. (2022). Two-stage anaerobic digestion: State of technology and perspective roles in future energy systems. *Bioresource Technology*, 360, 127633. <https://doi.org/10.1016/j.biortech.2022.127633>

- Ioannidou, S. M., Stylianou, E., Pateraki, C., Kookos, I., Rabaey, K., Koutinas, A., & Ladakis, D. (2023). Techno-economic and environmental sustainability assessment of succinic acid production from municipal biowaste using an electrochemical membrane bioreactor. *Chemical Engineering Journal*, 473, 145070. <https://doi.org/10.1016/j.cej.2023.145070>
- Jang, Y. S., Im, J. A., Choi, S. Y., Im Lee, J., & Lee, S. Y. (2014). Metabolic engineering of *Clostridium acetobutylicum* for butyric acid production with high butyric acid selectivity. *Metabolic Engineering*, 23, 165–174. <https://doi.org/10.1016/j.ymben.2014.03.004>
- Kaza, S., Yao, L., Bhada-Tata, P., & Van Woerden, F. (2018). *What a waste 2.0: A global snapshot of solid waste management to 2050*. World Bank Publications. <https://doi.org/10.1596/978-1-4648-1329-0>
- Kim, J. Y., Lee, J. A., Kim, G. B., Lee, Y., & Lee, S. Y. (2025). Succinic acid production by engineered *Mannheimia succiniciproducens* and its use in chemoenzymatic poly (butylene succinate) synthesis. *Biotechnology and Bioengineering*, 122(12), 3393–3405. <https://doi.org/10.1002/bit.70072>
- Lithourgidis, A. A., Kotsopoulos, T. A., Kalamaras, S. D., Skiadas, I. V., Kuglarz, M., Vigato, F., Alvarado-Morales, M., & Angelidaki, I. (2023). Bio-succinic acid production, up to pilot scale, by fermentation of industrial candy waste with *Actinobacillus succinogenes* 130Z and its downstream purification process. *Journal of Environmental Chemical Engineering*, 11(5), 110920. <https://doi.org/10.1016/j.jece.2023.110920>
- Malubhoy, Z., Bahia, F. M., de Valk, S. C., de Hulster, E., Rendulić, T., Ortiz, J. P. R., Xiberras, J., Klein, M., Mans, R., & Nevoigt, E. (2022). Carbon dioxide fixation via production of succinic acid from glycerol in engineered *Saccharomyces cerevisiae*. *Microbial Cell Factories*, 21(1), 102. <https://doi.org/10.1186/s12934-022-01817-1>
- Mitrea, L., Teleky, B.-E., Nemes, S.-A., Plamada, D., Varvara, R.-A., Pascuta, M.-S., Ciont, C., Cocean, A.-M., Medeleanu, M., Nistor, A., Rotar, A.-M., Pop, C.-R., & Vodnar, D.-C. (2024). Succinic acid – A run-through of the latest perspectives of production from renewable biomass. *Heliyon*, 10(3), e25551. <https://doi.org/10.1016/j.heliyon.2024.e25551>
- Mukherjee, P., Pal, S., & Sivaprakasam, S. (2025). Optimization of D-lactic acid biosynthesis from diverse carbon sources in mutant *Lactobacillus delbrueckii* subsp. *bulgaricus* via random mutagenesis. *Systems*

- Microbiology and Biomanufacturing*, 5(2), 702–721. <https://doi.org/10.1007/s43393-024-00316-1>
- Pamueangmun, P., Abdullahi, A. D., Kabir, M. H., Unban, K., Kanpiengjai, A., Venus, J., Shetty, K., Saenjum, C., & Khanongnuch, C. (2023). Lignocellulose degrading *Weizmannia coagulans* capable of enantiomeric L-lactic acid production via consolidated bioprocessing. *Fermentation*, 9(8), 761. <https://doi.org/10.3390/fermentation9080761>
- Panesar, P. S., Kennedy, J. F., Knill, C. J., & Kosseva, M. (2010). Production of L (+) lactic acid using *Lactobacillus casei* from whey. *Brazilian Archives of Biology and Technology*, 53(1), 219–226. <https://doi.org/10.1590/S1516-89132010000100027>
- Pu, Y., Tang, J., Wang, Q., & Abomohra, A. (2026). High-value caproate synthesis from food waste: Mechanisms, regulation, and sustainable scaling—A comprehensive review. *Journal of Environmental Management*, 402, 129138. <https://doi.org/10.1016/j.jenvman.2026.129138>
- Rajendran, N., & Han, J. (2023). Techno-economic analysis and life cycle assessment of poly (butylene succinate) production using food waste. *Waste Management*, 156, 168–176. <https://doi.org/10.1016/j.wasman.2022.11.037>
- Reddy, M. V., Mohan, S. V., & Chang, Y. C. (2018). Medium-chain fatty acids (MCFA) production through anaerobic fermentation using *Clostridium kluyveri*: Effect of ethanol and acetate. *Applied Biochemistry and Biotechnology*, 185(3), 594–605. <https://doi.org/10.1007/s12010-017-2674-2>
- Rochón, E., Banchieri, C., Barcos, M., Mihalik, J., & Lareo, C. (2024). Succinic acid production from sorghum grain: A comparative study of *Actinobacillus succinogenes* DSM 22257 and *Basfia succiniciproducens* DSM 22022 strains. *Biomass and Bioenergy*, 188, 107333. <https://doi.org/10.1016/j.biombioe.2024.107333>
- Sabbe, K., Boon, N., & Ganigué, R. (2025). Impact of temperature on butyric and caproic acid production in lactic acid chain elongation. *Journal of Environmental Chemical Engineering*, 13(5), 119152. <https://doi.org/10.1016/j.jece.2025.119152>
- Song, L., Cai, C., Chen, Z., Lin, C., Lv, Y., Ye, X., Liu, Y., Dai, X., & Liu, M. (2024). Lactic acid production from food waste: Advances in microbial fermentation and separation technologies. *Bioresource Technology*, 414, 131635. <https://doi.org/10.1016/j.biortech.2024.131635>

- Strazzera, G., Battista, F., Garcia, N. H., Frison, N., & Bolzonella, D. (2018). Volatile fatty acids production from food wastes for biorefinery platforms: A review. *Journal of Environmental Management*, 226, 278–288. <https://doi.org/10.1016/j.jenvman.2018.08.039>
- Sudhakar, M. P., & Dharani, G. (2022). Evaluation of seaweed for the production of lactic acid by fermentation using *Lactobacillus plantarum*. *Bioresource Technology Reports*, 17, 100890. <https://doi.org/10.1016/j.biteb.2021.100890>
- Swetha, T. A., Ananthi, V., Bora, A., Sengottuvelan, N., Ponnuchamy, K., Muthusamy, G., & Arun, A. (2023). A review on biodegradable polylactic acid (PLA) production from fermentative food waste – Its applications and degradation. *International Journal of Biological Macromolecules*, 234, 123703. <https://doi.org/10.1016/j.ijbiomac.2023.123703>
- Tang, J., Hu, Z., Pu, Y., Wang, X. C., & Abomohra, A. (2024). Bioprocesses for lactic acid production from organic wastes toward industrialization – A critical review. *Journal of Environmental Management*, 369, 122372. <https://doi.org/10.1016/j.jenvman.2024.122372>
- Ulčar, B., Regueira, A., Podojsteršek, M., Boon, N., & Ganigué, R. (2024). Why do lactic acid bacteria thrive in chain elongation microbiomes? *Frontiers in Bioengineering and Biotechnology*, 11, 1291007. <https://doi.org/10.3389/fbioe.2023.1291007>
- Vital, M., Howe, A. C., & Tiedje, J. M. (2014). Revealing the bacterial butyrate synthesis pathways by analyzing (meta)genomic data. *mBio*, 5(2), e00889-14. <https://doi.org/10.1128/mbio.00889-14>
- Vlaeminck, E., Uitterhaegen, E., Quataert, K., Delmulle, T., De Winter, K., & Soetaert, W. K. (2022). Industrial side streams as sustainable substrates for microbial production of poly(3-hydroxybutyrate) (PHB). *World Journal of Microbiology and Biotechnology*, 38(12), 238. <https://doi.org/10.1007/s11274-022-03416-z>
- Wang, H., Zhou, W., Gao, J., Ren, C., & Xu, Y. (2022). Revealing the characteristics of glucose- and lactate-based chain elongation for caproate production by *Caproicibacterium lactatifermentans* through transcriptomic, bioenergetic, and regulatory analyses. *mSystems*, 7(5), e00534-00522. <https://doi.org/10.1128/msystems.00534-22>
- Wang, Q., Yang, N., Cai, Y., Zhang, R., Wu, Y., Ma, W., Fu, C., Zhang, P., & Zhang, G. (2023). Advances in understanding entire process of medium chain carboxylic acid production from organic wastes via

- chain elongation. *Chemosphere*, 339, 139723.
<https://doi.org/10.1016/j.chemosphere.2023.139723>
- Wu, K.-K., Xu, P.-P., Zhao, L., Ren, N.-Q., & Zhang, Y.-F. (2024). Microbial conversion of carbon dioxide into premium medium-chain fatty acids: The progress, challenges, and prospects. *npj Materials Sustainability*, 2(1), 4.
<https://doi.org/10.1038/s44296-024-00008-w>
- Wu, S.-L., Long, Y., Wei, W., Shi, X., Shen, D., & Ni, B.-J. (2023). Co-electron donors driven medium-chain fatty acids production: Roles of electron donors, reaction kinetics and metabolic pathways. *Chemosphere*, 338, 139515.
<https://doi.org/10.1016/j.chemosphere.2023.139515>
- Yang, Y., Zhao, J., Zhou, Y., Xu, S., Ren, X., & Wei, Q. (2024). Progress on production of succinic acid by *Actinobacillus succinogenes* – New opportunities for cheap biomass and waste gas utilization. *Journal of Cleaner Production*, 434, 140005.
<https://doi.org/10.1016/j.jclepro.2023.140005>
- Yi, J., Choi, S., Han, M.-S., Lee, J. W., & Lee, S. Y. (2013). Production of succinic acid from renewable resources. In Yang, S.-T., El-Enshasy, H. A., & Thongchul, N. (Eds.), *Bioprocessing technologies in biorefinery for sustainable production of fuels, chemicals, and polymers* (pp. 317–330). John Wiley & Sons.
<https://doi.org/10.1002/9781118642047.ch17>
- Zhu, X., Zhou, Y., Wang, Y., Wu, T., Li, X., Li, D., & Tao, Y. (2017). Production of high-concentration n-caproic acid from lactate through fermentation using a newly isolated Ruminococcaceae bacterium CPB6. *Biotechnology for Biofuels*, 10(1), 102. <https://doi.org/10.1186/s13068-017-0788-y>
- Zhu, Y., Romain, C., & Williams, C. K. (2016). Sustainable polymers from renewable resources. *Nature*, 540(7633), 354–362.
<https://doi.org/10.1038/nature21001>